

# PALM BEACH ORTHOPAEDIC INSTITUTE, P.A.

G. CLAY BAYNHAM, M.D. | MICHAEL COONEY, M.D. | MELISA ESTES, M.D. | VINCENT FOWBLE, M.D. | KENNETH GERSZBERG, M.D.  
BRIAN HILL, M.D. | JOHN HINSON, M.D. | JAMES KERPSACK, M.D. | JUSTIN KEARSE, M.D. | MICHAEL LEIGHTON, M.D. | FERNANDO MORA, M.D.  
ANDREW NOBLE, M.D. | SCOTT NORRIS, D.O. | JOHN SCHILERO, D.P.M. | ANDREW SELTZER, D.O. | GARY WEXLER, M.D.

---

## **Rehabilitation Protocol** **Rotator Cuff Repair (Massive)**

### **PHASE I: Protected ROM** (*8 weeks*)

- May remove dressing and shower postop day # 3.
- Sutures are all underneath the skin and will dissolve on their own.
- Ice or cold flow systems encouraged for the first week at a minimum: should be used 3-4 times per day.
- Sling should be in place when not performing exercises.
- Hold pendulum exercises until 4 weeks following surgery
- May start active scapular mobility exercises at 6 weeks – Must keep the shoulder musculature relaxed.
- Avoid all active and active assistive exercises until cleared by the surgeon. This includes pulley exercises, wand and supine assisted exercises.
- Initiate exercise program 3 times per day:
  - Immediate elbow, forearm and hand range of motion out of sling
  - Passive external rotation of the shoulder to tolerance - instruct family member (start at 6 weeks)
  - PROM into scapular plane elevation to 120 degrees (start at 6 weeks)

### **PHASE II: Progressive ROM** (*8 to 12 weeks*)

- May discontinue sling.
- Lifting restriction of 5 pounds should be reinforced with patient.
- Start AAROM and AROM – includes pulleys, wand and supine gravity assisted exercises. Emphasize all motions except IR behind the back until 3 months.
- Isolate and strengthen scapular stabilizers.
- Progress PROM and capsular stretching of the shoulder as needed – elevation and external rotation as tolerated
- Avoid resisted training or strengthening. Avoid AROM in positions of subacromial impingement.

### **PHASE III:** (*>12 weeks*)

- Discontinue formal lifting restrictions.
- Start progressive rotator cuff and shoulder strengthening at 3 months – isometrics initially
- Initiate isotonic strengthening at 4 months (Theraband, dumbbells, Hughston's exercises, etc). Include home cuff strengthening program. Continue to emphasize scapular stabilizers.
- Equate active and passive range of motion. Encourage scapulohumeral mechanics during active shoulder motion.
- Simulate work/recreational activities as rotator cuff strength and endurance improve.

---

(561) 694-7776 | [www.pboi.com](http://www.pboi.com)

4215 BURNS RD, SUITE 100, PALM BEACH GARDENS, FL 33410 | 2055 MILITARY TR, SUITE 200, JUPITER, FL 33458  
1411 FLAGLER DR, SUITE 9800, WEST PALM BEACH, FL 33401 | 7701 SOUTHERN BLVD, SUITE 100, WEST PALM BEACH, FL 33411